

EGFR Mutation and Response of Lung Cancer to Gefitinib

TO THE EDITOR: Kobayashi et al. (Feb. 24 issue)¹ report that a second mutation in the gene encoding the epidermal growth factor receptor (*EGFR*), one resulting in a threonine-to-methionine substitution at amino acid position 790 (T790M), was associated with acquired resistance to gefitinib in their patient and that this mutant gene had been absent from the primary non-small-cell lung cancer. In a reanalysis of the data from the 397 subjects we have previously described,^{2,3} we identified two women who had never smoked who had non-small-cell lung cancer and harbored two *EGFR* mutations — T790M and a leucine-to-arginine substitution at amino acid position 858 (L858R) — in resected tumor specimens before treatment with chemotherapy or radiotherapy. Both patients later had recurrent disease and eventually died — outcomes suggesting that tumors with both the L858R and T790M mutations are very aggressive. One patient was treated with gefitinib and had progression.

These findings indicate the existence of cases with inherent double mutations and provide evidence that the T790M mutant genotype is an important factor conferring resistance to gefitinib in non-small-cell lung cancers containing *EGFR* sensitivity mutations. In addition, detecting T790M may be useful for predicting pretreatment resistance to *EGFR* tyrosine kinase inhibitors. Our observation, together with data from recent reports,^{1,4} may help clarify the role of *EGFR* mutations in the development of *EGFR*-related non-small-cell lung cancer and help establish effective strategies against specific subtypes of non-small-cell lung cancer.

Shinichi Toyooka, M.D.

Katsuyuki Kiura, M.D.

Okayama University Graduate School of Medicine and Dentistry
Okayama 700-8558, Japan
s_toyooka@nigeka2.hospital.okayama-u.ac.jp

Tetsuya Mitsudomi, M.D.

Aichi Cancer Center Hospital
Nagoya 464-8681, Japan

1. Kobayashi S, Boggon TJ, Dayaram T, et al. *EGFR* mutation and resistance of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2005;352:786-92.
2. Kosaka T, Yatabe Y, Endoh H, Kuwano H, Takahashi T, Mitsudomi T. Mutations of the epidermal growth factor receptor gene in lung cancer: biological and clinical implications. *Cancer Res* 2004;64:8919-23.

3. Tokumo M, Toyooka S, Kiura K, et al. The relationship between epidermal growth factor receptor mutations and clinicopathologic features in non-small cell lung cancers. *Clin Cancer Res* 2005;11:1167-73.

4. Pao W, Miller VA, Politi KA, et al. Acquired resistance of lung adenocarcinomas to gefitinib or erlotinib is associated with a second mutation in the *EGFR* kinase domain. *PLoS Med* 2005;2(3):e73.

THE AUTHORS REPLY: Dr. Toyooka and colleagues describe two patients whose lung tumors harbored a T790M mutation before treatment with chemotherapy or radiotherapy was begun and suggest that this mutation might be a marker of tumor aggressiveness as well as resistance to gefitinib therapy. In the cases we and others¹ have described, the T790M mutation was not found in specimens from untreated patients. Nevertheless, the possibilities do exist that this second mutation might be present in some tumors at a low frequency at the time of diagnosis and that tumor cells harboring the mutation might be enriched over time during treatment with gefitinib or erlotinib. By analogy, imatinib-resistant *BCR-ABL* mutations have, on occasion, been detected in specimens from patients with untreated chronic myeloid leukemia.^{2,3} We agree that such interesting findings should motivate further research to improve our understanding of the role of *EGFR* in non-small-cell lung cancers, to encourage the development of alternative *EGFR* inhibitors able to overcome such resistance mutations, and to incorporate the knowledge gained into clinical treatment.

Susumu Kobayashi, M.D., Ph.D.

Daniel G. Tenen, M.D.

Beth Israel Deaconess Medical Center
Boston, MA 02115

Balázs Halmos, M.D.

Case Western Reserve University
Cleveland, OH 44106

1. Pao W, Miller VA, Politi KA, et al. Acquired resistance of lung adenocarcinomas to gefitinib or erlotinib is associated with a second mutation in the *EGFR* kinase domain. *PLoS Med* 2005;2(3):e73.
2. Roche-Lestienne C, Soenen-Cornu V, Grardel-Duflos N, et al. Several types of mutations of the *Abl* gene can be found in chronic myeloid leukemia patients resistant to STI571, and they can pre-exist to the onset of treatment. *Blood* 2002;100:1014-8.
3. Roche-Lestienne C, Lai JL, Darre S, Facon T, Preudhomme C. A mutation conferring resistance to imatinib at the time of diagnosis of chronic myelogenous leukemia. *N Engl J Med* 2003;348:2265-6.